

## SITE SYNOPSIS

**SAC Site Name :** **LOWER RIVER SUIR** Candidate Special Area of Conservation (cSAC)

**SAC Site Code :** 002137

This site consists of the freshwater stretches of the River Suir immediately south of Thurles, the tidal stretches as far as the confluence with the Barrow/Nore immediately east of Cheekpoint in Co. Waterford and many tributaries including the Clodiagh in Co. Waterford, the Lingaun, Anner, Nier, Tar, Aherlow, Multeen and Clodiagh in Co. Tipperary. The Suir and its tributaries flows through the counties of Tipperary, Kilkenny and Waterford. Upstream of Waterford city, the swinging meanders of the Suir crisscross the Devonian sandstone rim of hard rocks no less than three times as they leave the limestone-floored downfold below Carrick. In the vicinity of Carrick-on-Suir the river follows the limestone floor of the Carrick Syncline. Upstream of Clonmel the river and its tributaries traverse Upper Palaeozoic Rocks, mainly the Lower Carboniferous Visean and Tournaisian. The freshwater stretches of the Clodiagh River in Co. Waterford traverse Silurian rocks, through narrow bands of Old Red Sandstone and Lower Avonian Shales before reaching the carboniferous limestone close to its confluence with the Suir. The Aherlow River flows through a Carboniferous limestone valley, with outcrops of Old Red Sandstone forming the Galtee Mountains to the south and the Slievenamuck range to the north. Glacial deposits of sands and gravels are common along the valley bottom, flanking the present-day river course.

The site is a candidate SAC selected for the presence of the priority habitats on Annex I of the E.U. Habitats Directive - alluvial wet woodlands and Yew Wood. The site is also selected as a candidate SAC for floating river vegetation, Atlantic salt meadows, Mediterranean salt meadows, old oak woodlands and eutrophic tall herbs, all habitats listed on Annex I of the E.U. Habitats Directive. The site is also selected for the following species listed on Annex II of the same directive - Sea Lamprey, River Lamprey, Brook Lamprey, Freshwater Pearl Mussel, Crayfish, Twaite Shad, Atlantic Salmon and Otter.

Alluvial wet woodland is declining habitat in Europe as a result of drainage and reclamation. The best examples of this type of woodland in the site are found on the islands just below Carrick-on-Suir and at Fiddown Island. Species occurring here include Almond Willow (*Salix triandra*), White Willow (*S. alba*), Grey Willow (*S. cinerea*), Osier (*S. viminalis*), with Iris (*Iris pseudacorus*), Hemlock Water-dropwort (*Oenanthe crocata*), Angelica (*Angelica sylvestris*), Pendulous Sedge (*Carex pendula*), Meadowsweet (*Filipendula ulmaria*) and Valerian (*Valeriana officinalis*). The terrain is littered with dead trunks and branches and intersected with small channels which carry small streams to the river. The bryophyte and lichen floras appear to be rich and require further investigation. A small plot is currently being coppiced and managed by National Parks and Wildlife. In the drier areas the wet woodland species merge with other tree and shrub species including Ash (*Fraxinus excelsior*), Hazel (*Corylus avellana*), Hawthorn (*Crataegus monogyna*) and Blackthorn (*Prunus spinosa*). This adds further to the ecological interest of this site.

Eutrophic tall herb vegetation occurs in association with the various areas of alluvial forest and elsewhere where the flood-plain of the river is intact. Characteristic species of the habitat include Meadowsweet (*Filipendula ulmaria*), Purple Loosestrife (*Lythrum salicaria*), Marsh Ragwort (*Senecio aquaticus*), Ground Ivy (*Glechoma hederacea*) and Hedge Bindweed (*Calystegia sepium*).

Old oak woodlands are also of importance at the site. The best examples are seen in Portlaw Wood which lies on both sides of the Clodiagh River. On the south-facing side the stand is more open and the Oaks (mainly *Quercus robur*) are well grown and spreading. Ivy (*Hedera helix*) and Bramble (*Rubus fruticosus*) are common on the ground, indicating relatively high light conditions. Oak regeneration is dense, varying in age from 0-40 years and Holly (*Ilex aquifolium*) is fairly common but mostly quite young. Across the valley, by contrast, the trees are much more closely spaced and though taller are poorly grown on average. There are no clearings; large Oaks extend to the boundary wall. In the darker conditions, Ivy is much rarer and Holly much more frequent, forming a closed canopy in places. Oak regeneration is uncommon since there are as yet few natural clearings. The shallowness of the soil on the north-facing slope probably contributes to the poor tree growth there. The acid nature of the substrate has induced a “mountain” type Oakwood community to develop. There is an extensive species list present throughout including an abundance of mosses, liverworts and lichens. The rare lichen *Lobaria pulmonaria*, an indicator of ancient woodlands, is found.

Inchinquilib Wood consists of three small separate sloping blocks of woodland in a valley cut by the young Multeen River and its tributaries through acidic Old Red Sandstone, and Silurian rocks. Two blocks, both with an eastern aspect, located to the north of the road, are predominantly of Sessile oak (*Quercus petraea*) and Hazel, with Downy Birch (*Betula pubescens*), Ash and Holly. The ground flora is quite mixed with for example Wood sedge (*Carex sylvatica*), Bluebell (*Hyacinthoides non-scriptus*), Primrose (*Primula vulgaris*), Wood-sorrel (*Oxalis acetosella*), Pignut (*Conopodium majus*) and Hard fern (*Blechnum spicant*). The base poor nature of the underlying rock is, to some extent masked by the overlying drift. The third block, to the south of the road, and with a northern aspect, is a similar although less mature mixture of Sessile Oak, Birch and Holly, the influence of the drift is more marked, with the occurrence of Wood anemone (*Anemone nemorosa*) amongst the ground flora.

Floating river vegetation is evident in the freshwater stretches of the River Suir and along many of its tributaries. Typical species found include Canadian Pondweed (*Elodea canadensis*), Milfoil (*Myriophyllum* spp.), Fennel Pondweed (*Potamogeton pectinatus*), Curled Pondweed (*P. crispus*), Perfoliate Pondweed (*P. perfoliatus*), Pond Water-crowfoot (*Ranunculus peltatus*), other Crowfoots (*Ranunculus* spp.) and the moss *Fontinalis antipyretica*. At a couple of locations along the river, Opposite-leaved Pondweed (*Groenlandia densa*) occurs. This species is protected under the Flora (Protection) Order, 1999.

The Aherlow River is fast-flowing and mostly follows a natural unmodified river channel. Submerged vegetation includes the aquatic moss *Fontinalis antipyretica* and Stream Water-crowfoot (*Ranunculus peltatus*), while shallow areas support species such as Reed Canary-grass (*Phalaris arundinacea*), Brooklime (*Veronica beccabunga*) and Water Mint (*Mentha aquatica*). The river bank is fringed in places with Alder (*Alnus glutinosa*) and Willows (*Salix* spp.).

The Multeen River is fast flowing, mostly gravel-bottomed and appears to follow a natural unmodified river channel. Water Crowfoots occur in abundance and the aquatic moss *Fontinalis antipyretica* is also common. In sheltered shallows, species such as Water-cress (*Rorippa nasturtium-aquaticum*) and Water-starworts (*Callitriche* spp.) occur. The river channel is fringed for most of its length with Alder, Willow and a narrow strip of marshy vegetation.

Salt meadows occur below Waterford City in old meadows where the embankment is absent, or has been breached, and along the tidal stretches of some of the in-flowing rivers below Little Island.

There are very narrow, non-continuous bands of this habitat along both banks. More extensive areas are also seen along the south bank at Ballynakill, the east side of Little Island, and in three large salt meadows between Ballynakill and Cheekpoint. The Atlantic and Mediterranean sub types are generally intermixed. The species list is extensive and includes Red Fescue (*Festuca rubra*), Oraches (*Atriplex* spp.), Sea Aster (*Aster tripolium*), Sea Couch Grass (*Elymus pycnanthus*), frequent Sea Milkwort (*Glaux maritima*), occasional Wild Celery (*Apium graveolens*), Parsley Water-dropwort (*Oenanthe lachenalii*), English Scurvygrass (*Cochlearia anglica*) and Sea Arrowgrass (*Triglochin maritima*). These species are more representative of the Atlantic sub-type of the habitat. Common Cord-grass (*Spartina anglica*), is rather frequent along the main channel edge and up the internal channels. The legally protected (Flora (Protection) Order, 1999) Meadow Barley (*Hordeum secalinum*) grows at the landward transition of the saltmarsh. Sea Rush (*Juncus maritimus*), an indicator of the Mediterranean salt meadows, also occurs.

Other habitats at the site include wet and dry grassland, marsh, reed swamp, improved grassland, coniferous plantations, deciduous woodland, scrub, tidal river, stony shore and mudflats. The most dominant habitat adjoining the river is improved grassland, although there are wet fields with species such as Yellow Flag (*Iris pseudacorus*), Meadow Sweet (*Filipendula ulmaria*), Rushes (*Juncus* spp.), Meadow Buttercup (*Ranunculus acris*) and Cuckoo Flower (*Cardamine pratensis*).

Cabragh marshes, just below Thurles, lie in a low-lying tributary valley into which the main river floods in winter. Here there is an extensive area of Common Reed (*Phragmites australis*) with associated marshland and peaty fen. The transition between vegetation types is often well displayed. A number of wetland plants of interest occur, in particular the Narrow-leaved Bulrush (*Typha angustifolia*), Bottle Sedge (*Carex rostrata*) and Blunt-flowered Rush (*Juncus subnodulosus*). The marsh is naturally eutrophic but it has also the nutritional legacy of the former sugar factory which discharged into it through a number of holding lagoons, now removed. Production is high which is seen in the size of such species as Celery-leaved Buttercup (*Ranunculus sceleratus*) as well as in the reeds themselves.

Throughout the Lower River Suir site are small areas of woodland other than those described above. These tend to be a mixture of native and non-native species, although there are some areas of semi-natural wet woodland with species such as Ash and Willow. Cahir Park Woodlands is a narrow tract of mixed deciduous woodland lying on the flat-lying floodplain of the River Suir. This estate woodland was planted over one hundred years ago and it contains a large component of exotic tree species. However, due to original planting and natural regeneration there is now a good mix of native and exotic species. About 5km north west of Cashel, Ardmayle pond is a long, possibly artificial water body running parallel to the River Suir. It is partly shaded by planted Lime (*Tilia* hybrids), Sycamore (*Acer pseudoplatanus*) and the native Alder. Growing beneath the trees are shade tolerant species such as Remote sedge (*Carex remota*).

The site is of particular conservation interest for the presence of a number of Annex II animal species, including Freshwater Pearl Mussel (*Margaritifera margaritifera* and *M. m. durrovensis*), Freshwater Crayfish (*Austropotamobius pallipes*), Salmon (*Salmo salar*), Twaite Shad (*Alosa fallax fallax*), three species of Lampreys - Sea Lamprey (*Petromyzon marinus*), Brook Lamprey (*Lampetra planeri*) and River Lamprey (*Lampetra fluviatilis*) and Otter (*Lutra lutra*). This is one of only three known spawning grounds in the country for Twaite Shad.

The site also supports populations of several other animal species. Those which are listed in the Irish Red Data Book include Daubenton's Bat (*Myotis daubentoni*), Natterer's Bat (*M. nattereri*),

Pipistrelle (*Pipistrellus pipistrellus*), Pine Marten (*Martes martes*), Badger (*Meles meles*), the Irish Hare (*Lepus timidus hibernicus*), Smelt (*Osmerus eperlanus*) and the Frog (*Rana temporaria*). Breeding stocks of Carp are found in Kilsheelan Lake. This is one of only two lakes in the country which is known to have supported breeding Carp. Carp require unusually high summer water temperatures to breed in Ireland and the site may therefore support interesting invertebrate populations.

Parts of the site have also been identified as of ornithological importance for a number of Annex I (EU Birds Directive) bird species, including Greenland White-fronted Goose (10), Golden Plover (1490), Whooper Swan (7) and Kingfisher. Figures given in brackets are the average maximum counts from 4 count areas within the site for the three winters between 1994 and 1997. Wintering populations of migratory birds use the site. Flocks are seen in Coolfinn Marsh and also along the reedbeds and saltmarsh areas of the Suir. Coolfinn supports nationally important numbers of Greylag Geese on a regular basis. Numbers between 600 and 700 are recorded. Other species occurring include Mallard (21), Teal (159), Widgeon (26), Tufted Duck (60), Pintail (4), Pochard (2), Little Grebe (2), Black-tailed Godwit (20), Oystercatcher (16), Lapwing (993), Dunlin (101), Curlew (195), Redshank (28), Greenshank (4) and Green Sandpiper (1). Nationally important numbers of Lapwing (2750) were recorded at Faithlegg in the winter of 1996/97. In Cabragh marshes there is abundant food for surface feeding wildfowl which total at 1,000 or so in winter. Widgeon, Teal and Mallard are numerous and the latter has a large breeding population - with up to 400 in summer. In addition, less frequent species like Shoveler and Pintail occur and there are records for both Whooper and Bewick's swans. Kingfisher, a species that is listed on Annex I of the EU Birds Directive, occurs along some of the many tributaries throughout the site.

Landuse at the site consists mainly of agricultural activities including grazing, silage production, fertilising and land reclamation. The grassland is intensively managed and the rivers are therefore vulnerable to pollution from run-off of fertilisers and slurry. Arable crops are also grown. Fishing is a main tourist attraction on stretches of the Suir and some of its tributaries and there are a number of Angler Associations, some with a number of beats. Fishing stands and styles have been erected in places. Both commercial and leisure fishing takes place on the rivers. The Aherlow River is a designated Salmonid Water under the EU Freshwater Fish Directive. Other recreational activities such as boating, golfing and walking are also popular. Several industrial developments, which discharge into the river, border the site including three dairy related operations and a tannery.

The Lower River Suir contains excellent examples of a number of Annex I habitats, including the priority habitat Alluvial Forest. The site also supports populations of several Annex II animal species and a number of Red Data Book animal species. The presence of two legally protected plants (Flora (Protection) Order, 1999) and the ornithological importance of the river adds further to the ecological interest of this site.

National Parks & Wildlife Service, 17.05.2005

Source: National Parks & Wildlife Service website,  
<http://www.npws.ie/media/Media,4167,en.pdf>

## SITE SYNOPSIS

**SITE NAME:** CABRAGH WETLANDS Proposed National Heritage Area (pNHA)  
**SITE CODE:** 001934

There are two parts to this site, both situated close to the River Suir near Thurles. The largest section is the Cabragh marshes which lie in a low-lying tributary valley into which the main river floods in winter. Here there is an extensive area of Common Reed (*Phragmites australis*) with associated marshland and peaty fen. The transition between vegetation types is often well displayed. A number of wetland plants of interest occur, in particular the Narrow-leaved Bulrush (*Typha angustifolia*), BottleSedge (*Carex vesicaria*) and Blunt-flowered Rush (*Juncus subnodulosus*).

The site is naturally eutrophic, but it has also the nutritional legacy of the former sugar factory which discharged into it through a number of holding lagoons, now removed. Production is high which is seen in the size of such species as Celery-leaved Buttercup (*Ranunculus sceleratus*) as well as in the reeds themselves. There is abundant food for surface feeding wildfowl which total at 1,000 or so in winter. Widgeon, teal and mallard are numerous and the latter has a large breeding population- with up to 400 in summer. In addition, less frequent species like shoveler and pintail occur and there are records for both whooper and Bewick's swans. Old records, (e.g. of constant green sandpipers) include the sugar factory lagoons so have to be quoted with care.

The second site is the Tank wetland in Ardbaun, north of the town. This is higher up a tributary valley than Cabragh and is mainly fed by springs of lime-rich groundwater. The abundance of Saw Sedge (*Cladium mariscus*) in the reedbeds indicates this feature and charophyte algae are also likely to be important though they have not yet been examined. Smaller numbers of wildfowl occur here, especially since, unlike the Cabragh marshes, it is shot over in winter. The site is adjoined by a refuse dump so there is some local littering in evidence. There is no evidence of major pollution from leachate at present.

Source: National Parks & Wildlife Service,  
undated; [post-closure of sugar factory, but predates closure of Thurles landfill in 1999]



## SITE SYNOPSIS

**SITE NAME:** **KILLOUGH HILL** Proposed National Heritage Area (pNHA)  
**SITE CODE:** 000959

The site surrounds a mounting quarry between Thurles and Cashel. The quarry exploits a limestone escarpment which had an abrupt north slope formed by the termination of its beds while the other side is a gentle slope conforming more or less perfectly with the inclination of the beds. Rock is now exposed on top and on the south slope where it has been eroded into limestone pavement.

The southern side of the hill is partly under grass, and drift soil normally overlies the limestone. Ash is well established here with some elder and scattered oak trees but only small patches of closed canopy are found.

Higher up and to the north side a hazel wood becomes dominant with some ash and rowan (*Sobus auscuparia*) and shrubs such as *Euonymus europaeus* (spindle-tree), *Viburnum opulus* (guelder rose) and *Rosa canina* and *R. arvensis* (roses). The woodland floor has a characteristic assemblage of herbs and mosses, with abundant Oxalis (*Oxalis acetosella*) wood sorrel (*Endymion non-scriptus*) and bluebell, and the woodland has a typical range of invertebrates.

The limestone grassland is dominated by Sheep's fescue (*Festuca ovina*) with characteristic lime loving plants, such as Lady's bedstraw (*Galium nevur*) and Centre thistle (*Centaurea vulgaris*). The pavement area is vegetated by a characteristic mixture of variously adapted plant species.

The woodland on Killough Hill is of relatively recent origin but is developing well with a very good ground flora. It also houses a large bird population and varied mammals. The open area of pavement and limestone grassland are perhaps more important since they are a rare feature in the country. To date only four exposures of pavement are known outside the western Clare-Galway area.

16 February, 1995

Source: National Parks & Wildlife Service.



## SITE SYNOPSIS

**SITE NAME:**           **LAFFANSBRIDGE** Proposed National Heritage Area (pNHA)

**SITE CODE:** 000965

This site, which comprises five separate areas, is situated close to the village of Laffansbridge which lies c.3km west of Killenaule and 13km east-north-east of Cashel. The site is underlain by Carboniferous limestone rocks and shallow calcium-rich soils.

The main vegetation type present at the site is calcareous grassland; this includes such plant species as Quaking Grass (*Briza media*), Downy Oat-grass (*Avenula pubescens*), Ladies Bedstraw (*Galium verum*) and Bulbous Buttercup (*Ranunculus bulbosus*).

While this herb-rich sward is of conservation interest itself, being a threatened habitat throughout Europe, Laffansbridge is of particular significance for the large population of Green-winged Orchid (*Orchis morio*), a nationally scarce species, that it supports. In places the population at this site is substantial, such as west of the railway at Manserghshill where some 6000 flowering spikes of this species have been recorded.

Limestone outcrops have prevented machinery access and preserved these unimproved grassland fragments from intensive fertilisation. However, they remain vulnerable to other developments such as quarrying and housing, which have damaged similar areas outside of the site. They are also sensitive to grazing management; low levels of grazing in some areas has allowed the establishment and spread of Gorse (*Ulex europaeus*), to the detriment of the herb-rich grassland.

8.4.2005

Source:           National Parks & Wildlife Service.